



Isolation and Genomic Analysis of an Intracellular *Mycobacterium gordonae* from a Free-Living *Acanthamoeba* sp. in a Hospital Environment in Lima, Peru

Alfonso M. Cabello-Vilchez,^{a,b} Alejandra Dávila-Barclay,^c  Pablo Tsukayama^{a,c,d}

^aInstituto de Medicina Tropical Alexander von Humboldt, Universidad Peruana Cayetano Heredia, Lima, Peru

^bLaboratorio de Protistas Patógenos, Universidad Privada Norbert Wiener, Lima, Peru

^cLaboratorio de Genómica Microbiana, Facultad de Ciencias y Filosofía, Universidad Peruana Cayetano Heredia, Lima, Peru

^dParasites and Microbes Programme, Wellcome Sanger Institute, Hinxton, United Kingdom

ABSTRACT *Mycobacterium gordonae* is a nontuberculous mycobacterium found in diverse environments and is considered an opportunistic pathogen in immunocompromised patients. We report the draft genome sequence of a *Mycobacterium gordonae* strain isolated from a free-living amoeba found in a nosocomial environment in Lima, Peru.

Mycobacterium *gordonae* is an opportunistic nontuberculous mycobacterium (NTM) ubiquitous in water systems (1–4), where it shares a habitat with free-living amoebae (FLA) (5–8). Infections are associated with immunocompromised patients but may occur in immunocompetent hosts (9–15). Exposure to water networks and bodies poses a potential risk for disease (16, 17).

Previous reports have shown that NTM can associate with amoeba hosts in the environment (5, 8, 18). *Mycobacterium gordonae* isolate NTM676 was found intracellularly in an amoebal cyst previously isolated from a faucet surface biofilm in a tertiary-level hospital in Lima, Peru. After 48 h of incubation at room temperature, a positive *Acanthamoeba* sp. culture grown in nonnutritive agar with a lawn of *Escherichia coli* ATCC 25922 was observed under phase-contrast microscopy (Fig. 1A). Page's amoeba saline solution was added to the plate for 1 h and then streaked into a 15-mL conical tube. The centrifuged sediment was placed in a new tube and left at room temperature for 30 days. Aliquots were taken weekly for Ziehl-Neelsen staining.

We observed a slow-growing intracellular acid-fast bacillus (AFB) at week 4 inside amoebal trophozoites and cysts (Fig. 1B). A gentamicin and streptomycin solution (300 μ g/mL) was added to the tube and incubated at room temperature until trophozoites were no longer present (Fig. 1C). The remaining cysts containing AFB were cultured in nonnutritive agar until trophozoite transformation and then lysed with sodium dodecyl sulfate (SDS) and decontaminated using modified Petroff's method (19). The sediment was washed with phosphate buffer and inoculated in Löwenstein-Jensen (LJ) medium at 30°C for 25 days. A single small smooth orange scotochromogenic colony suggested the growth of an NTM-like organism (Fig. 1D). Further screenings were not conducted prior to whole-genome sequencing.

DNA was extracted from colonies restreaked in LJ medium (Fig. 1E) using the GeneJET DNA purification kit (Thermo Fisher Scientific, Waltham, MA). Genomic libraries were prepared using the Nextera XT kit (Illumina, San Diego, CA) and sequenced on an Illumina MiSeq instrument and a V2 kit, generating 2,082,926 250-bp paired-end raw sequences. Reads were quality checked and trimmed with FastQC v0.11.9 (20) and Trimmomatic v0.39 (21), removing reads shorter than 20 bp. *De novo* assembly using SPAdes v3.15.4 (22) resulted in 314 contigs, a total length of 7,266,850 bp with an N_{50} of 42,384 bp, mean coverage of 51 \times , and a GC content of 66.2%.

Editor Steven R. Gill, University of Rochester School of Medicine and Dentistry

Copyright © 2022 Cabello-Vilchez et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Pablo Tsukayama, pablo.tsukayama@upch.pe.

The authors declare no conflict of interest.

Received 3 August 2022

Accepted 1 October 2022

Published 17 October 2022




FIG 1 (A) Acanthamoeba trophozoites (phase-contrast microscopy, $\times 60$ magnification). (B) Acanthamoeba cysts with acid-fast bacillus microcolonies, Ziehl-Neelsen stain (light microscopy, $\times 40$ magnification). (C) Positive Ziehl-Neelsen-stained bacillus (light microscopy, $\times 100$ magnification). (D) Single orange nontuberculous mycobacterial colony in Löwenstein-Jensen medium. (E) Restreaked mycobacterial colony in Löwenstein-Jensen medium.

Resistome prediction with CARD 3.2.3 RGI 5.2.1 (23) identified genes associated with resistance to fosfomycin (*murA*, mutation C117D), rifamycin (*rbpA*; *rpoB*, mutations D516G, H526T, L511R), and macrolides (*mtrA*) with a $>90\%$ identity threshold and aminoglycosides [*aac(2)-Ic*] with an $>80\%$ identity threshold. PathogenFinder v1.1 (24) indicated a 75% probability of being a human pathogen, matching with 25 known virulence protein families, including ESAT-6 and others related to the ESX-5 secretion system, with a median identity threshold of 93.5%. All programs were run on default parameters.

This draft genome sequence reveals the antibiotic resistance and pathogenic potential of a nontuberculous mycobacterium found endosymbiotically in a free-living amoeba from hospital water systems.

Data availability. The whole-genome shotgun project for *M. gordonae* NTM676 is available at DDBJ/ENA/GenBank under the accession number [JANFXG010000000](https://doi.org/10.1038/s41579-020-0331-1). Raw sequence reads are available in the Sequence Read Archive (SRA) under the accession number [SRR14802570](https://doi.org/10.1038/s41579-020-0331-1).

ACKNOWLEDGMENTS

This study was supported by CONCYTEC/ProCiencia grant 011-2019.

We thank Instituto de Medicina Tropical Alexander von Humboldt and Hospital Nacional Cayetano Heredia for enabling sampling of hospital surfaces and Diego Cuicapuza for curating the QC/assembly pipeline. A.M.C.V. thanks S.I.R.A.

REFERENCES

- Johansen MD, Herrmann JL, Kremer L. 2020. Non-tuberculous mycobacteria and the rise of *Mycobacterium abscessus*. *Nat Rev Microbiol* 18:392–407. <https://doi.org/10.1038/s41579-020-0331-1>.
- Honda JR, Viridi R, Chan ED. 2018. Global environmental nontuberculous mycobacteria and their contemporaneous man-made and natural niches. *Front Microbiol* 9:2029. <https://doi.org/10.3389/fmicb.2018.02029>.

3. Cabello-Vílchez AM, Núñez-Ato RG. 2018. Aislamiento y caracterización molecular de micobacterias no tuberculosas en el sistema de distribución de agua en un hospital de Lima (Perú). *Rev Biosalud* 17:7–24.
4. Zulu M, Monde N, Nkhoma P, Malama S, Munyeme M. 2021. Nontuberculous mycobacteria in humans, animals, and water in Zambia: a systematic review. *Front Trop Dis* 2:679501. <https://doi.org/10.3389/ftd.2021.679501>.
5. Ovrutsky AR, Chan ED, Kartalija M, Bai X, Jackson M, Gibbs S, Falkinham JO, Iseman MD, Reynolds PR, McDonnell G, Thomas V. 2013. Cooccurrence of free-living amoebae and nontuberculous mycobacteria in hospital water networks, and preferential growth of mycobacterium avium in *Acanthamoeba lenticulata*. *Appl Environ Microbiol* 79:3185–3192. <https://doi.org/10.1128/AEM.03823-12>.
6. Amisshah NA, Gryseels S, Tobias NJ, Ravadgar B, Suzuki M, Vandellannoote K, Durnez L, Leirs H, Stinear TP, Portaels F, Ablordey A, Eddyani M. 2014. Investigating the role of free-living amoebae as a reservoir for Mycobacterium ulcerans. *PLoS Negl Trop Dis* 8:e3148. <https://doi.org/10.1371/journal.pntd.0003148>.
7. Delafont V, Mougari F, Cambau E, Joyeux M, Bouchon D, Hécharde Y, Moulin L. 2014. First evidence of amoebae-mycobacteria association in drinking water network. *Environ Sci Technol* 48:11872–11882. <https://doi.org/10.1021/es5036255>.
8. Thomas V, McDonnell G. 2007. Relationship between mycobacteria and amoebae: ecological and epidemiological concerns. *Let Appl Microbiol* 45:349–357. <https://doi.org/10.1111/j.1472-765X.2007.02206.x>.
9. Jarikre LN. 1992. Mycobacterium gordonae genitourinary disease. *Genitourin Med* 68:45–46. <https://doi.org/10.1136/sti.68.1.45>.
10. Foti C, Sforza V, Rizzo C, De Pascale G, Bonamonte D, Conserva A, Tarantino A, Stella C, Cantore S, Grassi RF, Ballini A, De Vito D, Angelini G. 2009. Cutaneous manifestations of Mycobacterium gordonae infection described for the first time in Italy: a case report. *Cases J* 2:6828. <https://doi.org/10.4076/1757-1626-2-6828>.
11. Chen Y, Jiang J, Jiang H, Chen J, Wang X, Liu W, Chen Z, Shi Y, Zhang W, Wang H. 2017. Mycobacterium gordonae in patient with facial ulcers, nosebleeds, and positive T-SPOT.TB test result, China. *Emerg Infect Dis* 23:1204–1206. <https://doi.org/10.3201/eid2307.162033>.
12. Pamparino S, Valente I, Tagliafico A, Dentone C, Bassetti M, Mennella S, Calabrese M, Garlaschi A. 2020. A very rare case of Mycobacterium gordonae infection of the breast. *Breast J* 26:2229–2232. <https://doi.org/10.1111/tbj.14086>.
13. Ustinova V, Smirnova T, Blagodatikh K, Varlamov D, Sochivko D, Larionova E, Andreevskaya S, Andrievskaya I, Chernousova L. 2016. First draft genome sequence of a Mycobacterium gordonae clinical isolate. *Genome Announc* 4:6–7. <https://doi.org/10.1128/genomeA.00638-16>.
14. Weinberger M, Berg SL, Feuerstein IM, Pizzo PA, Witebsky FG. 1992. Disseminated infection with Mycobacterium gordonae: report of a case and critical review of the literature. *Clin Infect Dis* 14:1229–1239. <https://doi.org/10.1093/clinids/14.6.1229>.
15. Mazumder SA, Hicks A, Norwood J. 2010. Mycobacterium gordonae pulmonary infection in an immunocompetent adult. *N Am J Med Sci* 2: 205–207.
16. Thomson R, Tolson C, Carter R, Coulter C, Huygens F, Hargreaves M. 2013. Isolation of nontuberculous mycobacteria (NTM) from household water and shower aerosols in patients with pulmonary disease caused by NTM. *J Clin Microbiol* 51:3006–3011. <https://doi.org/10.1128/JCM.00899-13>.
17. Oriani A, Sierra F, Baldini M. 2018. Effect of chlorine on Mycobacterium gordonae and Mycobacterium chubuense in planktonic and biofilm state. *Int J Mycobacteriol* 7:122–127. https://doi.org/10.4103/ijmy.ijmy_30_18.
18. Gryseels S, Amisshah D, Durnez L, Vandellannoote K, Leirs H, de Jonckheere J, Silva MT, Portaels F, Ablordey A, Eddyani M. 2012. Amoebae as potential environmental hosts for Mycobacterium ulcerans and other mycobacteria, but doubtful actors in Buruli ulcer epidemiology. *PLoS Negl Trop Dis* 6:e1764. <https://doi.org/10.1371/journal.pntd.0001764>.
19. Kent P, Kubica G. 1985. Public health mycobacteriology: a guide for the level III laboratory. Centers for Disease Control, Atlanta, GA.
20. Andrews S. 2010. FastQC: a quality control tool for high throughput sequence data. <https://www.bioinformatics.babraham.ac.uk/projects/fastqc/>. Accessed 30 November 2019.
21. Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30:2114–2120. <https://doi.org/10.1093/bioinformatics/btu170>.
22. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
23. Alcock BP, Raphenya AR, Lau TT, Tsang KK, Bouchard E, Edalatmand A, Huynh W, Nguyen A-LV, Cheng AA, Liu S, Min SY, Miroshnichenko A, Tran H-K, Werfalli RE, Nasir JA, Oloni M, Speicher DJ, Florescu A, Singh B, Faltyn M, Hernandez-Koutouchcheva A, Sharma AN, Bordeleau E, Pawlowski AC, Zubyk HL, Dooley D, Griffiths E, Maguire F, Winsor GL, Beiko RG, Brinkman FS, Hsiao WW, Domselaar GV, McArthur AG. 2020. CARD 2020: antibiotic resistance surveillance with the Comprehensive Antibiotic Resistance Database. *Nucleic Acids Res* 48:D517–D525. <https://doi.org/10.1093/nar/gkz935>.
24. Cosentino S, Voldby Larsen M, Møller Aarestrup F, Lund O. 2013. PathogenFinder—distinguishing friend from foe using bacterial whole genome sequence data. *PLoS One* 8:e77302. <https://doi.org/10.1371/journal.pone.0077302>.